

PostScript

LETTERS TO MP

If you have a burning desire to respond to a paper published in *Molecular Pathology*, why not make use of our "rapid response" option?

Log on to our website (www.molpath.com), find the paper that interests you, and send your response via email by clicking on the "eLetters" option in the box at the top right hand corner.

Providing it isn't libellous or obscene, it will be posted within seven days. You can retrieve it by clicking on "read eLetters" on our homepage.

The editors will decide as before whether to also publish it in a future paper issue.

The assessment of DNA and RNA preservation in liquid based cytology media

The Papanicolaou cervical smear method has long been used for cervical cancer screening, but there is a trend to develop liquid based cytology media and monolayer techniques to reduce false negative rates. These last two types of technique seem to increase the sensitivity of cervical cancer detection,¹ and also provide cellular residues that are useful either for spotting additional slices or performing molecular techniques, such as hybrid capture or the polymerase chain reaction (PCR).² Different media exist and to date the PreservCyt™ medium used in the ThinPrep® Pap Test™ method (Cytoc, Boxborough, Massachusetts, USA) has been validated by the Food

and Drug Administration as an alternative method to the conventional smear. Different methods of human papillomavirus (HPV) detection have been compared and the PCR seems to have higher analytical sensitivity than hybrid capture, at least on the samples processed with the PreservCyt liquid.² A previous report, which tested serial dilutions of an HPV positive cell line, showed that the PreservCyt liquid is suitable for molecular techniques because of good preservation of total nucleic acids (RNA and DNA).³

We compared two different media—PreservCyt and Seroa® (Seroa, Monaco)⁴—used for cervical cytology for DNA and RNA based techniques; in particular, for cellular residues (that is, in routine practice). We randomly selected 10 cellular residues of cervical smears fixed in each medium and performed DNA and RNA extraction with protocols that have already been published.⁵ Briefly, after centrifugation, the pellets were split into two aliquots—one for DNA and one for RNA extraction. The extraction provided equal amounts of DNA in the two series (ranges, 200–1000 ng/µl). However, the amounts of RNA were low, with no clear differences between the two media, varying from 0 to 15 ng/µl. To compare the quality of the DNA and RNA samples, we performed PCR techniques with primers specific for the β actin gene β1/β2 (PCR products: DNA, 560 bp; RNA, 360 bp) after 15, 20, 25, and 35 cycles.⁵ PCR amplification of DNA was efficient in all cases (with primers specific for β actin or c-raf-1), and a visible band was obtained after 15 cycles, with slight differences of staining intensity by ethidium bromide. Regarding RNA preservation, six of the 20 samples gave rise to visible reverse transcription PCR (RT-PCR) products (PreservCyt, n = 3 and Seroa, n = 3). The best signal (plateau) was achieved after 35 cycles. RNA amplification failure is probably the result of poor RNA

preservation, rather than differences in the amounts of cell residue, which were similar in the two media. This is confirmed by the DNA amplification results where both media gave similar results. Our data indicate that in cellular residues of routinely processed samples both media are satisfactory and equivalent for DNA based techniques (PCR, hybrid capture), but the amount of RNA is frequently insufficient even for highly sensitive RT-PCR techniques. Because RNA degradation occurs frequently in routine conditions, the addition of RNase inhibitors to the media should be investigated as a means of circumventing this problem.

P Trempat, R C Zenou, P Brousset

Department of Pathology, UPCM UPR2163 CNRS, CHU Purpan, Avenue de Grande Bretagne, 31059 Toulouse Cedex; brousset.p@chu-toulouse.fr

References

- 1 **Monsonogo J**, Autillo-Touati A, Bergeron C, *et al*. Liquid-based cytology for primary cervical cancer screening: a multi-centre study. *Br J Cancer* 2001;**84**:360–6.
- 2 **Cope JU**, Hildesheim A, Schiffman MH, *et al*. Comparison of the hybrid capture tube test and PCR for detection of human papillomavirus DNA in cervical specimens. *J Clin Microbiol* 1997;**35**:2262–5.
- 3 **Tarkowski TA**, Rajeevan MS, Lee DR, *et al*. Improved detection of viral RNA isolated from liquid-based cytology samples. *Mol Diagn* 2001;**6**:125–30.
- 4 **Zerat L**. Le frottis cervical en suspension liquide. *Revue du Praticien Gynecologie Obstetrique* 1999;**38**:19–22.
- 5 **Galoin S**, Daste G, Apoil PA, *et al*. Polymerase chain reaction on cerebrospinal fluid cells in the detection of leptomeningeal involvement by B-cell lymphoma and leukaemia: a novel strategy and its implications. *Br J Haematol* 1997;**99**:122–30.



The assessment of DNA and RNA preservation in liquid based cytology media

P Trempat, R C Zenou and P Brousset

Mol Path 2002 55: 125

doi: 10.1136/mp.55.2.125

Updated information and services can be found at:

<http://mp.bmj.com/content/55/2/125.full.html>

References

These include:

This article cites 4 articles, 1 of which can be accessed free at:

<http://mp.bmj.com/content/55/2/125.full.html#ref-list-1>

Article cited in:

<http://mp.bmj.com/content/55/2/125.full.html#related-urls>

Email alerting service

Receive free email alerts when new articles cite this article. Sign up in the box at the top right corner of the online article.

Notes

To request permissions go to:

<http://group.bmj.com/group/rights-licensing/permissions>

To order reprints go to:

<http://journals.bmj.com/cgi/reprintform>

To subscribe to BMJ go to:

<http://group.bmj.com/subscribe/>